

A Handbook of Statistical Analyses Using R

Brian S. Everitt and Torsten Hothorn



Conditional Inference: Guessing Lengths, Suicides, Gastrointestinal Damage, and Newborn Infants

3.1 Introduction

3.2 Conditional Test Procedures

3.3 Analysis Using R

3.3.1 Estimating the Width of a Room Revised

The unconditional analysis of the room width estimated by two groups of students in Chapter ?? lead to the conclusion that the estimates in metres are slightly larger than the estimates in feet. Here, we reanalyse these data in a conditional framework. First, we convert metres into feet and store the vector of observations in a variable y :

```
R> data("roomwidth", package = "HSAUR")
R> convert <- ifelse(roomwidth$unit == "feet", 1, 3.28)
R> feet <- roomwidth$unit == "feet"
R> metre <- !feet
R> y <- roomwidth$width * convert
```

The test statistic is simply the difference in means

```
R> T <- mean(y[feet]) - mean(y[metre])
R> T
```

```
[1] -8.858893
```

In order to approximate the conditional distribution of the test statistic T we compute 9999 test statistics for shuffled y values. A permutation of the y vector can be obtained from the `sample` function.

```
R> meandiffs <- double(9999)
R> for (i in 1:length(meandiffs)) {
+   sy <- sample(y)
+   meandiffs[i] <- mean(sy[feet]) - mean(sy[metre])
+ }
```

The distribution of the test statistic T under the null hypothesis of independence of room width estimates and groups is depicted in Figure 3.1. Now, the value of the test statistic T for the original unshuffled data can be compared

```
R> hist(meandiffs)
R> abline(v = T, lty = 2)
R> abline(v = -T, lty = 2)
```




Figure 3.1 Approximated conditional distribution of the difference of mean `roomwidth` estimates in the feet and metres group under the null hypothesis. The vertical lines show the negative and positive absolute value of the test statistic T obtained from the original data.

with the distribution of T under the null hypothesis (the vertical lines in Figure 3.1). The p -value, i.e., the proportion of test statistics T larger than 8.859 or smaller than -8.859 is

```
R> greater <- abs(meandiffs) > abs(T)
R> mean(greater)
```

```
[1] 0.00740074
```

with a confidence interval of

```
R> binom.test(sum(greater), length(greater))$conf.int
```

```
[1] 0.005815511 0.009282160
attr(,"conf.level")
[1] 0.95
```

Note that the approximated conditional p -value is roughly the same as the p -value reported by the t -test in Chapter 2.

```
R> library("coin")
R> independence_test(y ~ unit, data = roomwidth,
+                   distribution = exact())
      Exact General Independence Test

data: y by unit (feet, metres)
Z = -2.5491, p-value = 0.008492
alternative hypothesis: two.sided
```

Figure 3.2 R output of the exact permutation test applied to the roomwidth data.

```
R> wilcox_test(y ~ unit, data = roomwidth,
+             distribution = exact())
      Exact Wilcoxon-Mann-Whitney Test

data: y by unit (feet, metres)
Z = -2.1981, p-value = 0.02763
alternative hypothesis: true mu is not equal to 0
```

Figure 3.3 R output of the exact conditional Wilcoxon rank sum test applied to the roomwidth data.

3.3.2 Crowds and Threatened Suicide

```
R> data("suicides", package = "HSAUR")
R> fisher.test(suicides)
      Fisher's Exact Test for Count Data

data: suicides
p-value = 0.0805
alternative hypothesis: true odds ratio is not equal to 1
95 percent confidence interval:
 0.7306872 91.0288231
sample estimates:
odds ratio
 6.302622
```

Figure 3.4 R output of Fisher's exact test for the suicides data.

3.3.3 Gastrointestinal Damages

Here we are interested in the comparison of two groups of patients, where one group received a placebo and the other one Misoprostol. In the trials shown here, the response variable is measured on an ordered scale – see Table ?? . Data from four clinical studies are available and thus the observations are naturally grouped together. From the *data.frame* `Lanza` we can construct a three-way table as follows:

```
R> data("Lanza", package = "HSAUR")
R> xtabs(~ treatment + classification + study, data = Lanza)
, , study = I
```

	classification				
treatment	1	2	3	4	5
Misoprostol	21	2	4	2	0
Placebo	2	2	4	9	13

```
, , study = II
```

	classification				
treatment	1	2	3	4	5
Misoprostol	20	4	6	0	0
Placebo	8	4	9	4	5

```
, , study = III
```

	classification				
treatment	1	2	3	4	5
Misoprostol	20	4	3	1	2
Placebo	0	2	5	5	17

```
, , study = IV
```

	classification				
treatment	1	2	3	4	5
Misoprostol	1	4	5	0	0
Placebo	0	0	0	4	6

For the first study, the null hypothesis of independence of treatment and gastrointestinal damage, i.e., of no treatment effect of Misoprostol, is tested by

```
R> library("coin")
R> cmh_test(classification ~ treatment, data = Lanza,
+           scores = list(classification = c(0, 1, 6, 17, 30)),
+           subset = Lanza$study == "I")
```

Asymptotic Linear-by-Linear Association Test

```
data: classification (ordered) by treatment (Misoprostol, Placebo)
```

```
Z = -5.371, p-value = 7.83e-08
alternative hypothesis: two.sided
```

and, by default, the conditional distribution is approximated by the corresponding limiting distribution. The p -value indicates a strong treatment effect. For the second study, the asymptotic p -value is a little bit larger

```
R> cmh_test(classification ~ treatment, data = Lanza,
+           scores = list(classification = c(0, 1, 6, 17, 30)),
+           subset = Lanza$study == "II")
```

Asymptotic Linear-by-Linear Association Test

```
data: classification (ordered) by treatment (Misoprostol, Placebo)
Z = -3.4733, p-value = 0.000514
alternative hypothesis: two.sided
```

and we make sure that the implied decision is correct by calculating a confidence interval for the exact p -value

```
R> p <- cmh_test(classification ~ treatment, data = Lanza,
+               scores = list(classification = c(0, 1, 6, 17, 30)),
+               subset = Lanza$study == "II", distribution =
+               approximate(B = 19999))
```

```
R> pvalue(p)
```

```
[1] <5.00025e-05
99 percent confidence interval:
0.000000000 0.000264894
```

The third and fourth study indicate a strong treatment effect as well

```
R> cmh_test(classification ~ treatment, data = Lanza,
+           scores = list(classification = c(0, 1, 6, 17, 30)),
+           subset = Lanza$study == "III")
```

Asymptotic Linear-by-Linear Association Test

```
data: classification (ordered) by treatment (Misoprostol, Placebo)
Z = -5.3065, p-value = 1.118e-07
alternative hypothesis: two.sided
```

```
R> cmh_test(classification ~ treatment, data = Lanza,
+           scores = list(classification = c(0, 1, 6, 17, 30)),
+           subset = Lanza$study == "IV")
```

Asymptotic Linear-by-Linear Association Test

```
data: classification (ordered) by treatment (Misoprostol, Placebo)
Z = -3.9675, p-value = 7.262e-05
alternative hypothesis: two.sided
```

At the end, a separate analysis for each study is unsatisfactory. Because the design of the four studies is the same, we can use `study` as a block variable and perform a global linear-association test investigating the treatment effect

of Misoprostol in all four studies. The block variable can be incorporated into the *formula* by the `|` symbol.

```
R> cmh_test(classification ~ treatment | study, data = Lanza,
+           scores = list(classification = c(0, 1, 6, 17, 30)))
```

Asymptotic Linear-by-Linear Association Test

```
data:  classification (ordered) by
       treatment (Misoprostol, Placebo)
       stratified by study
Z = -9.1443, p-value < 2.2e-16
alternative hypothesis: two.sided
```

Based on this result, a strong treatment effect can be established.

3.3.4 Teratogenesis

In this example, the medical doctor (MD) and the research assistant (RA) assessed the number of anomalies (0, 1, 2 or 3) for each of 395 babies:

```
R> anomalies <- c(235, 23, 3, 0, 41, 35, 8, 0,
+               20, 11, 11, 1, 2, 1, 3, 1)
R> anomalies <- as.table(matrix(anomalies,
+                               ncol = 4, dimnames = list(MD = 0:3, RA = 0:3)))
R> anomalies
```

	RA			
MD	0	1	2	3
0	235	41	20	2
1	23	35	11	1
2	3	8	11	3
3	0	0	1	1

We are interested in testing whether the number of anomalies assessed by the medical doctor differs structurally from the number reported by the research assistant. Because we compare *paired* observations, i.e., one pair of measurements for each newborn, a test of marginal homogeneity (a generalisation of McNemar's test, see Chapter 2) needs to be applied:


```
R> mh_test(anomalies)
```

```
Asymptotic Marginal Homogeneity Test
```

```
data: response by  
      conditions (MD, RA)  
      stratified by block  
chi-squared = 21.227, df = 3, p-value = 9.446e-05
```

The p -value indicates a deviation from the null hypothesis. However, the levels of the response are not treated as ordered. Similar to the analysis of the gastrointestinal damage data above, we can take this information into account by the definition of an appropriate score. Here, the number of anomalies is a natural choice:

```
R> mh_test(anomalies, scores = list(response = c(0, 1, 2, 3)))
```

```
Asymptotic Marginal Homogeneity Test for Ordered Data
```

```
data: response (ordered) by  
      conditions (MD, RA)  
      stratified by block  
Z = -4.5847, p-value = 4.545e-06  
alternative hypothesis: two.sided
```

In our case, one can conclude that the assessment of the number of anomalies differs between the medical doctor and the research assistant.